



INFANT FORMULAS: Medical Request Required

Provided by the Wisconsin WIC Program to Partially Breastfed and Fully Formula Fed Infants (Birth to 12 Months of Age)

A qualifying medical condition, is required using the Wisconsin WIC Request for Medical Formula/Food – Infants and Children form and is subject to WIC RD approval.

Formula	Use: Information provided is from manufacturers' website and materials. Manufacturers frequently alter the ingredients in the formula products, which may affect the nutrient content. For the most current and accurate information, refer to the product label and/or manufacturer website.	Protein Sources	CHO Sources	Fat Sources	Product Form, Size, Amount	By Age, Maximum Number Cans Per Month and Prepared Ounces Per Day					
						0-3 Mo		4-5 Mo		6-11 Mo ¹	
						cans	oz	cans	oz	cans	oz
Similac NeoSure Abbott Nutrition	Use: A high-calorie, infant formula for premature infants in need of extra calories and nutrients. For use as a preterm, post discharge formula. Contains higher levels of protein, vitamins, and minerals than standard term formulas to promote greater weight gain and growth. Kosher, Halal. Standard Dilution: 22 kcal/fl oz, 1 scoop to 2 oz water Caution: For use only under medical supervision.	(11% kcals) 50:50 ³ Nonfat milk, whey protein concentrate	(40% kcal) Corn syrup solids, lactose	(50% kcals) Soy oil, high oleic safflower oil, coconut oil, 25% MCT, DHA	Powder ² 13.1 oz can 87 fl oz	10	29	11	32	8	23
					Ready to Feed ² 32 fl oz bottle	26	28	28	30	20	21
Enfamil NeuroPro EnfaCare Mead Johnson	Use: A milk-based 22 kcal discharge formula designed for premature infants who have left the hospital but still require the extra calories, nutrients, and minerals of a specialized infant formula. Can be used through the first year of life. Contains higher levels of vitamins and minerals than standard term formulas. Standard Dilution: 22 kcal/fl oz, 1 scoop to 2 oz water Caution: For use only under medical supervision.	(11% kcals) 60:40 ³ whey protein concentrate, nonfat milk	(42% kcals) Corn syrup solids, lactose; <i>Ready to Feed: contains maltodextrin</i>	(47% kcals) High oleic vegetable oil (sunflower and/or safflower oil) soy oil, 22% MCT, coconut oil, DHA, ARA	Powder ² 12.8 oz can 82 fl oz	10	27	11	30	8	22
Enfamil AR Mead Johnson	Use: Term infant milk-based formula with rice starch (formula thickens in acid environment of stomach), Kosher. Standard Dilution: 20 kcal/fl oz, 1 scoop to 2 oz water Caution: Not for infants born prematurely.	(10% kcals) 20:80 ³ Nonfat milk	(44% kcals) Rice starch, lactose, maltodextrin	(46% kcals) Vegetable oil (palm olein, soy, coconut, and high oleic sunflower), DHA, ARA	Powder ² 12.9 oz can 91 fl oz	9	28	10	31	7	22
					Ready to Feed ² 6 pk of 8 oz containers 48 fl oz	17 6pks	28	19 6pks	30	13 6pks	22
Extensive HA Gerber	Use: Hypoallergenic infant formula for the dietary management of cow's milk protein allergy. With probiotic <i>B. lactis</i> , lactose-free. Standard dilution: 20 kcal/fl oz, 1 scoop to 1 oz water Caution: For use only under medical supervision.	(10% kcals) Hydrolyzed whey protein isolate	(44% kcals) corn maltodextrin, potato starch	(46% kcals) 49% MCT, Vegetable oils (soy, high oleic sunflower and high 2-palmitic vegetable oil), DHA, ARA	Powder ² 14.1 oz can 96 fl oz	9	29	10	32	7	22

Formula	Use: Information provided is from manufacturers' website and materials. Manufacturers frequently alter the ingredients in the formula products, which may affect the nutrient content. For the most current and accurate information, refer to the product label and/or manufacturer website.	Protein Sources	CHO Sources	Fat Sources	Product Form, Size, Amount	By Age, Maximum Number Cans Per Month and Prepared Ounces Per Day					
						0-3 Mo		4-5 Mo		6-11 Mo ¹	
						cans	oz	cans	oz	cans	oz
Nutramigen w/Enflora LGG (powder) Mead Johnson	Use: Hypoallergenic infant formula designed for infants who are allergic to the intact protein in cow's milk and soy formulas. Contains extensively hydrolyzed casein protein to help avoid an immune system response by reducing allergen exposure, and the probiotic branded LGG to help support the strength of the intestinal barrier and support digestive health. Gluten-free, Lactose-free, galactose-free, sucrose-free. Standard dilution: 20 kcal/fl oz, 1 scoop to 2 oz water Caution: For use only under medical supervision. Not recommended for routine use in very low birth rate (VLBW) infants, as these infants may be at increased risk of developing gastrointestinal complications.	(11% kcals) Casein hydrolysate ⁴ , amino acids	(41% kcals) Corn syrup solids, modified corn starch	(48% kcals) Vegetable oils (palm olein, coconut, soy, high oleic sunflower), DHA, ARA	Powder ² 12.6 oz can 87 fl oz	10	29	11	32	8	23
Nutramigen (liquids) Mead Johnson	Use: Hypoallergenic formula for infants who are allergic to intact proteins in cow's milk and soy formulas. Contains extensively hydrolyzed protein and is proven effective for managing colic due to cow's milk allergy. Lactose-free Standard dilution: 20 kcal/fl oz, 1 scoop to 2 oz water Caution: For use only under medical supervision. Not recommended for routine use in VLBW infants, as these infants may be at increased risk of developing gastrointestinal complications.	(11% kcals) Casein hydrolysate ⁴ , amino acids	(41% kcals) Corn syrup solids, modified corn starch	(48% kcals) Vegetable oils (palm olein, soy, coconut, high oleic sunflower), DHA, ARA	Concentrate 13 fl oz can 26 fl oz	31	27	34	29	24	21
					Ready to Feed ² 32 fl oz bottle	26	28	28	30	20	21
Similac Alimentum Abbott Nutrition	Use: Hypoallergenic formula for infants with severe food allergies, sensitivities to intact protein, protein maldigestion, or fat malabsorption. Gluten-free, Lactose-free, corn-free (<i>Ready to Feed only</i>) Standard dilution: 20 kcal/fl oz, 1 scoop to 2 oz water Caution: For use only under medical supervision.	(11% kcals) Casein hydrolysate ⁴ , amino acids	(41% kcals) Corn maltodextrin, sucrose (<i>Powder</i>)	(48% kcals) Safflower oil, 33% MCT, soy oil, DHA, ARA	Powder ² 12.1 oz can 87 fl oz	10	27	11	31	8	23
			Sucrose, modified tapioca starch (<i>RTF</i>)		Ready to Feed ² 32 fl oz bottle	26	28	28	30	20	21
Pregestimil Mead Johnson	Use: Hypoallergenic infant formula designed for infants who experience fat malabsorption and who may also be sensitive to intact proteins. Fat malabsorption or steatorrhea may be associated with cystic fibrosis, short bowel syndrome, intractable diarrhea, and severe protein calorie malnutrition. Gluten-free, lactose-free, galactose-free, sucrose-free. Standard dilution: 20 kcal/fl oz, 1 scoop to 2 oz water Caution: For use only under medical supervision. Not recommended for routine use in VLBW infants who may be at increased risk of developing gastrointestinal complications.	(11% kcals) Casein hydrolysate ⁴ , amino acids	(41% kcals) Corn syrup solids, dextrose, modified corn starch	(48% kcals) 55% MCT oil, soy oil, corn oil, high oleic safflower in powder, DHA, ARA	Powder ² 16 oz can 112 fl oz	7-8	27	8	30	6	22

Formula	Use: Information provided is from manufacturers' website and materials. Manufacturers frequently alter the ingredients in the formula products, which may affect the nutrient content. For the most current and accurate information, refer to the product label and/or manufacturer website.	Protein Sources	CHO Sources	Fat Sources	Product Form, Size, Amount	By Age, Maximum Number Cans Per Month and Prepared Ounces Per Day					
						0-3 Mo		4-5 Mo		6-11 Mo ¹	
						cans	oz	cans	oz	cans	oz
Neocate Infant DHA & ARA Nutricia	Use: An elemental, hypoallergenic formula for infants with cow and soy milk allergy, multiple food protein intolerance, short bowel syndrome, eosinophilic esophagitis, gastroesophageal reflux, and other medical conditions that require an amino-acid based diet. Dairy-free, soy oil-free, sucrose-free. Kosher Pareve, Halal. Standard dilution: 20 kcal/fl oz 1 scoop to 1 oz water Contraindications: For use only under medical supervision.	(11.2% kcals) Free amino acids	(43.1% kcals) Corn syrup solids	(45.7% kcals) Refined vegetable oil (33% MCT, high oleic sunflower oil, sunflower oil, canola oil) DHA, ARA	Powder ² 14.1 oz can 97 fl oz	8-9	29	9-10	32	7	26
EleCare DHA & ARA Infant Abbott	Use: An amino acid-based formula for infants who cannot tolerate intact or hydrolyzed protein. It is indicated for the dietary management of protein maldigestion, malabsorption, severe food allergies, short-bowel syndrome, eosinophilic GI disorders, GI-tract impairment, or other conditions in which an amino acid-based diet is required. Lactose-free, galactose-free, fructose-free, milk protein-free, soy protein-free, gluten-free. Halal. Standard dilution: 20 kcal/fl oz 1 scoop to 2 oz water Caution: For use only under medical supervision.	(15% kcals) 100% free amino acids	(42% kcals) Corn syrup solids	(43% kcals) High oleic safflower oil, 33% MCT, soy oil, DHA, ARA	Powder ² 14.1 oz can 95 fl oz	9	29	10	32	7	22
Similac PM 60/40 Abbott	Use: Infant formula for infants and children with impaired renal function or who would benefit from lower mineral intake. Calcium-to-phosphorus ration and content designed to treat serum calcium disorders (hypercalcemia and hypocalcemia due to hyperphosphatemia). Gluten-free. Kosher, Halal. Standard dilution: 20 kcal/fl oz 1 scoop to 2 oz water Caution: For use only under medical supervision. Additional iron should be supplied from other sources. In conditions where the infant is losing abnormal quantities of one or more electrolytes, it may be necessary to supply electrolytes from sources other than the formula. It may be necessary to supply low-birth-weight infants weighing less than 1500 g at birth additional calcium, phosphorus, and sodium during periods of rapid growth.	(9% kcals) Whey protein concentrate and sodium caseinate	(41% kcal) Lactose	(50% kcal) High oleic safflower, soy, and coconut oils	Powder ² 14.1 oz can 102 fl oz	8	30	9	34	6-7	23

¹ Infants 6-11 months of age may receive the same monthly maximum as infants 4 through 5 months of age when infant foods are contraindicated for medical reasons.
² Infant cases with serious infections secondary to *Enterobacter sakazakii* have been reported in the literature that were traced back to powdered infant formulas contaminated with this organism. Preterm infants and those with underlying medical conditions may be at higher risk of developing infection. FDA recommends sterile, commercial Ready-to-Feed (RTF) formulas for immunocompromised infants including preterm infants. WIC routinely provides powdered formula. However, RTF exempt formulas may be provided when the local WIC Project RD verifies that the water supply is unsafe to drink, if the caregiver has difficulty mixing powder or concentrate properly, if RTF better accommodates the participant's condition (e.g., immunocompromised), or if RTF improves the participant's compliance in consuming the prescribed formula.
³ Human milk is estimated to be 40:60 casein to whey, but because cow's milk whey is primarily lactoglobulin and human milk whey is primarily lactalbumin, an advantage of a particular casein:whey ratio has not been established.
⁴ Not Kosher/Halal since pork enzyme used to hydrolyze protein.