

NATALITY

| | | | | | | | | | | | |
|--|--------|-----------------------|--------------------|----------------------------------|---------|---|--------------------------|----------------|---------|----------------|--|
| TOTAL LIVE BIRTHS | 528 | Birth Order | | Births | Percent | Marital Status of Mother | | Births | Percent | | |
| Crude Live Birth Rate | 11.3 | First | | 204 | 39 | Married | | 345 | 65 | | |
| General Fertility Rate | 59.8 | Second | | 177 | 34 | Not married | | 183 | 35 | | |
| Live births with reported congenital anomalies | 6 1.1% | Third | | 85 | 16 | Unknown | | 0 | 0 | | |
| Delivery Type | Births | Percent | Fourth or higher | | 62 | 12 | Education of Mother | | Births | Percent | |
| Vaginal after previous cesarean | 10 | 2 | Unknown | | 0 | 0 | Elementary or less | | 29 | 5 | |
| Forceps | 3 | 1 | 1st Prenatal Visit | | Births | Percent | Some high school | | 58 | 11 | |
| Other vaginal | 374 | 71 | 1st trimester | | 438 | 83 | High school | | 180 | 34 | |
| Primary cesarean | 84 | 16 | 2nd trimester | | 71 | 13 | Some college | | 154 | 29 | |
| Repeat cesarean | 57 | 11 | 3rd trimester | | 13 | 2 | College graduate | | 107 | 20 | |
| Birthweight | Births | Percent | No visits | | 4 | 1 | Unknown | | . | . | |
| <1,500 gm | 7 | 1.3 | Prenatal Visits | | Births | Percent | Smoking Status of Mother | | Births | Percent | |
| 1,500-2,499 gm | 27 | 5.1 | No visits | | 4 | 1 | Smoker | | 99 | 19 | |
| 2,500+ gm | 494 | 93.6 | 1-4 | | 14 | 3 | Nonsmoker | | 429 | 81 | |
| Unknown | 0 | 0 | 5-9 | | 92 | 17 | Unknown | | 0 | 0 | |
| | | | 10-12 | | 278 | 53 | | | | | |
| | | | 13+ | | 136 | 26 | | | | | |
| | | | Unknown | | 4 | 1 | | | | | |
| ----- | | | | | | | | | | | |
| | | Race/Ethnicity | | Low Birthweight (under 2,500 gm) | | Trimester of First Prenatal Visit | | | | | |
| | | Births Percent | | Births Percent | | 1st Trimester | | 2nd Trimester | | Other/Unknown | |
| | | White | | 29 5.8 | | Births Percent | | Births Percent | | Births Percent | |
| | | Black/Afr.American | | . | | 417 84 | | 65 13 | | 17 3 | |
| | | American Indian | | 1 11.1 | | 5 63 | | 2 25 | | 1 13 | |
| | | Hispanic/Latino | | 2 22.2 | | 5 56 | | 3 33 | | 1 11 | |
| | | Laotian/Hmong | | . | | 8 89 | | 1 11 | | . | |
| | | Other/Unknown | | * . | | * . | | * . | | * . | |
| ----- | | | | | | | | | | | |
| | | Age of Mother | | Low Birthweight (under 2,500 gm) | | Trimester of First Prenatal Visit | | | | | |
| | | Births Fertility Rate | | Births Percent | | 1st Trimester | | 2nd Trimester | | Other/Unknown | |
| | | <15 | | . | | Births Percent | | Births Percent | | Births Percent | |
| | | 15-17 | | 2 18.2 | | 4 36 | | 5 45 | | 2 18 | |
| | | 18-19 | | 2 5.9 | | 30 88 | | 4 12 | | . | |
| | | 20-24 | | 13 7.6 | | 138 81 | | 27 16 | | 6 4 | |
| | | 25-29 | | 9 5.4 | | 146 87 | | 18 11 | | 4 2 | |
| | | 30-34 | | 6 6.2 | | 83 86 | | 11 11 | | 3 3 | |
| | | 35-39 | | 1 3.0 | | 27 82 | | 4 12 | | 2 6 | |
| | | 40+ | | 1 7.1 | | 10 71 | | 2 14 | | 2 14 | |
| | | Unknown | | . | | . | | . | | . | |
| | | Teenage Births | | 45 28 | | * Data not reported if age or race category has less than 5 births. | | | | | |

MORBIDITY

MORTALITY

REPORTED CASES OF SELECTED DISEASES*

| Disease | Number |
|-------------------------|--------|
| Campylobacter Enteritis | 8 |
| Giardiasis | 7 |
| Hepatitis Type A | 0 |
| Hepatitis Type B** | <5 |
| Hepatitis NANB/C | 11 |
| Legionnaire's | <5 |
| Lyme | 43 |
| Measles | 0 |
| Meningitis, Aseptic | <5 |
| Meningitis, Bacterial | 0 |
| Mumps | 0 |
| Pertussis | 21 |
| Salmonellosis | <5 |
| Shigellosis | 0 |
| Tuberculosis | <5 |

* 2005 provisional data.
 ** Includes all positive HBsAg test results.

| Sexually Transmitted Disease | Number |
|------------------------------|--------|
| Chlamydia Trachomatis | 67 |
| Genital Herpes | 14 |
| Gonorrhea | 6 |
| Syphilis | 0 |

IMMUNIZATIONS

| Children in Grades K-12 by Compliance Level | Number |
|---|--------|
| Compliant | 8,209 |
| Non-compliant | 35 |
| Percent Compliant | 99.6 |

Note: Death rates (except infant) are per 100,000 population. Infant, neonatal, and postneonatal death rates are per 1,000 live births. Perinatal and fetal death rates are per 1,000 live births plus fetal deaths. Rates are not calculated for fewer than 20 deaths.

| TOTAL DEATHS | | 487 |
|------------------|--------|--------|
| Crude Death Rate | | 1,046 |
| Age | Deaths | Rate |
| 1-4 | 1 | . |
| 5-14 | 1 | . |
| 15-19 | 3 | . |
| 20-34 | 10 | . |
| 35-54 | 35 | 253 |
| 55-64 | 36 | 645 |
| 65-74 | 87 | 2,346 |
| 75-84 | 131 | 4,839 |
| 85+ | 180 | 14,229 |

| Infant Mortality | | Deaths | Rate |
|------------------|---|--------|------|
| Total Infant | 3 | . | . |
| Neonatal | 2 | . | . |
| Postneonatal | 1 | . | . |
| Unknown | . | . | . |

| Race of Mother | | Deaths | Rate |
|----------------|---|--------|------|
| White | 3 | . | . |
| Black | . | . | . |
| Hispanic | . | . | . |
| Laotian/Hmong | . | . | . |
| Other/Unknown | . | . | . |

| Birthweight | | Deaths | Rate |
|----------------|---|--------|------|
| <1,500 gm | 2 | . | . |
| 1,500-2,499 gm | 1 | . | . |
| 2,500+ gm | . | . | . |
| Unknown | . | . | . |

| Perinatal Mortality | | Deaths | Rate |
|---------------------|---|--------|------|
| Total Perinatal | 4 | . | . |
| Neonatal | 2 | . | . |
| Fetal | 2 | . | . |

| Selected Underlying Cause | Deaths | Rate |
|---------------------------|--------|------|
| Heart Disease (total) | 138 | 296 |
| Ischemic heart disease | 95 | 204 |
| Cancer (total) | 108 | 232 |
| Trachea/Bronchus/Lung | 28 | 60 |
| Colorectal | 11 | . |
| Female Breast | 9 | .* |
| Cerebrovascular Disease | 40 | 86 |
| Lower Resp. Disease | 24 | 52 |
| Pneumonia & Influenza | 9 | . |
| Accidents | 23 | 49 |
| Motor vehicle | 10 | . |
| Diabetes | 8 | . |
| Infect./Parasitic Dis. | 11 | . |
| Suicide | 4 | . |

* Based on female population.

ALCOHOL AND DRUG ABUSE AS UNDERLYING OR CONTRIBUTING CAUSE OF DEATH

| Alcohol/Tobacco Use/Other Drugs | Deaths | Rate |
|---------------------------------|--------|------|
| Alcohol | 8 | . |
| Tobacco Use | 80 | 172 |
| Other Drugs | 3 | . |

MOTOR VEHICLE CRASHES

Note: These data are based on location of crash, not on residence.

| Type of Motor Vehicle Crash | Persons Injured | Persons Killed |
|-----------------------------|-----------------|----------------|
| Total Crashes | 437 | 14 |
| Alcohol-Related | 50 | 3 |
| With Citation: | | |
| For OWI | 31 | 0 |
| For Speeding | 14 | 0 |
| Motorcyclist | 26 | 1 |
| Bicyclist | 9 | 0 |
| Pedestrian | 5 | 0 |

2005 HOSPITALIZATIONS

| DISEASE / AGE GROUP | Number | Per 1,000 Pop | Average Stay (Days) | Average Charge | Charge Per Capita | DISEASE / AGE GROUP | Number | Per 1,000 Pop | Average Stay (Days) | Average Charge | Charge Per Capita |
|---|--------|---------------|---------------------|----------------|-------------------|--|--------|---------------|---------------------|----------------|-------------------|
| Injury-Related: All | | | | | | Alcohol-Related | | | | | |
| Total | 381 | 8.2 | 4.3 | \$16,099 | \$132 | Total | 104 | 2.2 | 4.1 | \$4,952 | \$11 |
| <18 | 26 | 2.6 | 1.7 | \$8,947 | \$23 | 18-44 | 59 | 3.7 | 3.9 | \$4,008 | \$15 |
| 18-44 | 89 | 5.6 | 3.7 | \$17,292 | \$96 | 45-64 | 40 | 3.1 | 4.2 | \$5,189 | \$16 |
| 45-64 | 86 | 6.7 | 3.8 | \$15,569 | \$104 | Pneumonia and Influenza | | | | | |
| 65+ | 180 | 23.4 | 5.3 | \$16,795 | \$394 | Total | 334 | 7.2 | 4.0 | \$8,749 | \$63 |
| Injury: Hip Fracture | | | | | | <18 | 45 | 4.5 | 1.9 | \$3,375 | \$15 |
| Total | 56 | 1.2 | 6.1 | \$16,901 | \$20 | 45-64 | 48 | 3.7 | 4.5 | \$12,688 | \$47 |
| 65+ | 47 | 6.1 | 6.4 | \$17,268 | \$106 | 65+ | 226 | 29.4 | 4.4 | \$9,233 | \$272 |
| Injury: Poisonings | | | | | | Cerebrovascular Disease | | | | | |
| Total | 22 | 0.5 | 1.6 | \$4,848 | \$2 | Total | 123 | 2.6 | 3.6 | \$12,193 | \$32 |
| 18-44 | 15 | . | . | . | . | 45-64 | 18 | . | . | . | . |
| Psychiatric | | | | | | 65+ | 100 | 13.0 | 3.6 | \$12,138 | \$158 |
| Total | 300 | 6.4 | 6.3 | \$6,193 | \$40 | Chronic Obstructive Pulmonary Disease | | | | | |
| <18 | 25 | 2.5 | 4.2 | \$3,778 | \$9 | Total | 101 | 2.2 | 3.4 | \$7,462 | \$16 |
| 18-44 | 173 | 10.8 | 5.7 | \$5,426 | \$59 | <18 | 9 | . | . | . | . |
| 45-64 | 73 | 5.7 | 9.1 | \$8,799 | \$50 | 18-44 | 4 | . | . | . | . |
| 65+ | 29 | 3.8 | 5.0 | \$6,297 | \$24 | 45-64 | 28 | 2.2 | 4.2 | \$8,839 | \$19 |
| Coronary Heart Disease | | | | | | 65+ | 60 | 7.8 | 3.5 | \$7,730 | \$60 |
| Total | 294 | 6.3 | 3.7 | \$30,041 | \$190 | Drug-Related | | | | | |
| 45-64 | 93 | 7.2 | 3.3 | \$34,696 | \$250 | Total | 44 | 0.9 | 10.3 | \$8,018 | \$8 |
| 65+ | 195 | 25.4 | 4.0 | \$28,151 | \$715 | 18-44 | 27 | 1.7 | 6.5 | \$5,470 | \$9 |
| Malignant Neoplasms (Cancers): All | | | | | | Total Hospitalizations | | | | | |
| Total | 178 | 3.8 | 5.3 | \$19,354 | \$74 | Total | 5,307 | 114.0 | 3.7 | \$11,104 | \$1,265 |
| 18-44 | 17 | . | . | . | . | <18 | 777 | 78.1 | 2.5 | \$3,563 | \$278 |
| 45-64 | 48 | 3.7 | 4.8 | \$18,989 | \$71 | 18-44 | 1,267 | 79.1 | 3.0 | \$7,339 | \$580 |
| 65+ | 113 | 14.7 | 5.4 | \$19,517 | \$287 | 45-64 | 1,050 | 81.3 | 4.2 | \$15,645 | \$1,272 |
| Neoplasms: Female Breast (rates for female population) | | | | | | 65+ | 2,213 | 288.1 | 4.4 | \$13,752 | \$3,962 |
| Total | 18 | . | . | . | . | ----- PREVENTABLE HOSPITALIZATIONS* ----- | | | | | |
| Neoplasms: Colo-rectal | | | | | | Total | 879 | 18.9 | 3.6 | \$8,126 | \$153 |
| Total | 25 | 0.5 | 7.4 | \$23,295 | \$13 | <18 | 83 | 8.3 | 1.7 | \$3,250 | \$27 |
| 65+ | 18 | . | . | . | . | 18-44 | 86 | 5.4 | 2.7 | \$7,852 | \$42 |
| Neoplasms: Lung | | | | | | 45-64 | 167 | 12.9 | 4.0 | \$10,629 | \$137 |
| Total | 10 | . | . | . | . | 65+ | 543 | 70.7 | 3.9 | \$8,144 | \$576 |
| Diabetes | | | | | | * Hospitalizations for conditions where timely and effective ambulatory care can reduce the likelihood of hospitalization. | | | | | |
| Total | 60 | 1.3 | 4.5 | \$14,436 | \$19 | | | | | | |
| 65+ | 31 | 4.0 | 5.2 | \$16,250 | \$66 | | | | | | |