

NATALITY

| TOTAL LIVE BIRTHS | | | Birth Order | | | Marital Status of Mother | | | | |
|--|--|--------|-------------|--------------------|------------------|--------------------------|---------|---------------------|--------------------------|---------|
| | | | Births | Percent | | | | | | |
| TOTAL LIVE BIRTHS | | | 234 | | First | 82 | 35 | Married | 161 | 69 |
| Crude Live Birth Rate | | | 9.3 | | Second | 70 | 30 | Not married | 73 | 31 |
| General Fertility Rate | | | 62.6 | | Third | 36 | 15 | Unknown | 0 | 0 |
| Live births with reported congenital anomalies | | | 7 | 3.0% | Fourth or higher | 46 | 20 | | | |
| | | | | | Unknown | 0 | 0 | Education of Mother | | |
| | | | | | | | | Births | Percent | |
| Delivery Type | | Births | Percent | 1st Prenatal Visit | | Births | Percent | Elementary or less | | |
| Vaginal after previous cesarean | | 7 | 3 | 1st trimester | | 183 | 78 | Some high school | | |
| Forceps | | 0 | 0 | 2nd trimester | | 39 | 17 | High school | | |
| Other vaginal | | 162 | 69 | 3rd trimester | | 11 | 5 | Some college | | |
| Primary cesarean | | 32 | 14 | No visits | | 0 | 0 | College graduate | | |
| Repeat cesarean | | 33 | 14 | Unknown | | 1 | <.5 | Unknown | | |
| | | | | | | | | | Smoking Status of Mother | |
| | | | | | | | | | Births | Percent |
| Birthweight | | Births | Percent | Prenatal Visits | | Births | Percent | of Mother | | |
| <1,500 gm | | 2 | 0.9 | No visits | | 0 | 0 | Smoker | | |
| 1,500-2,499 gm | | 15 | 6.4 | 1-4 | | 7 | 3 | Nonsmoker | | |
| 2,500+ gm | | 217 | 92.7 | 5-9 | | 59 | 25 | Unknown | | |
| Unknown | | 0 | 0 | 10-12 | | 92 | 39 | | | |
| | | | | 13+ | | 75 | 32 | | | |
| | | | | Unknown | | 1 | <.5 | | | |

| Race/Ethnicity | Births | | Low Birthweight (under 2,500 gm) | | Trimester of First Prenatal Visit | | | | | |
|--------------------|--------|---------|----------------------------------|---------|-----------------------------------|----|---------------|----|---------------|---|
| | Births | Percent | Births | Percent | 1st Trimester | | 2nd Trimester | | Other/Unknown | |
| White | 196 | 84 | 16 | 8.2 | 157 | 80 | 27 | 14 | 12 | 6 |
| Black/Afr.American | 2 | 1 | * | . | * | . | * | . | * | . |
| American Indian | 1 | <.5 | * | . | * | . | * | . | * | . |
| Hispanic/Latino | 34 | 15 | 1 | 2.9 | 22 | 65 | 12 | 35 | . | . |
| Asian | 1 | <.5 | * | . | * | . | * | . | * | . |
| Other/Unknown | 0 | 0 | . | . | . | . | . | . | . | . |

| Age of Mother | Fertility | | Low Birthweight (under 2,500 gm) | | Trimester of First Prenatal Visit | | | | | |
|---------------|-----------|------|----------------------------------|---------|-----------------------------------|----|---------------|----|---------------|----|
| | Births | Rate | Births | Percent | 1st Trimester | | 2nd Trimester | | Other/Unknown | |
| <15 | 0 | -- | . | . | . | . | . | . | . | . |
| 15-17 | 2 | 5 | * | . | * | . | * | . | * | . |
| 18-19 | 22 | 68 | 2 | 9.1 | 16 | 73 | 3 | 14 | 3 | 14 |
| 20-24 | 62 | 98 | 2 | 3.2 | 50 | 81 | 11 | 18 | 1 | 2 |
| 25-29 | 68 | 117 | 7 | 10.3 | 57 | 84 | 10 | 15 | 1 | 1 |
| 30-34 | 47 | 100 | 3 | 6.4 | 41 | 87 | 4 | 9 | 2 | 4 |
| 35-39 | 28 | 46 | 3 | 10.7 | 17 | 61 | 9 | 32 | 2 | 7 |
| 40+ | 5 | 7 | . | . | 2 | 40 | 1 | 20 | 2 | 40 |
| Unknown | 0 | -- | . | . | . | . | . | . | . | . |

| | | | | | | | | | | |
|----------------|----|----|---|--|--|--|--|--|--|--|
| Teenage Births | 24 | 34 | * Not reported if age or race category has fewer than 5 births. | | | | | | | |
|----------------|----|----|---|--|--|--|--|--|--|--|

MORBIDITY

MORTALITY

CONFIRMED CASES OF COMMUNICABLE DISEASES (2009)

| Disease | Number |
|---|--------|
| Campylobacter Enteritis | 7 |
| Giardiasis | <5 |
| Hepatitis Type A | 0 |
| Hepatitis Type B* | <5 |
| Hepatitis NANB/C | <5 |
| Legionnaire's | 0 |
| Lyme | 54 |
| Measles | 0 |
| N. Meningitis, Meningococcal | 0 |
| Meningitis, Bacterial | 0 |
| Mumps | 0 |
| Pertussis | <5 |
| Salmonellosis | <5 |
| Shigellosis | 7 |
| Tuberculosis | 0 |
| E-Coli 0157 | 0 |
| E-Coli, Non-0157 | 0 |
| Cryptosporidiosis | <5 |
| Streptococcus Pneum Invasive | <5 |
| All Streptococcal Diseases | <5 |
| Blastomycosis | 0 |
| Haemophilus flu, Invasive | 0 |
| Ehrlichiosis/Anaplasmosis | 0 |
| Influenza A-Novel | 22 |
| Kawasaki Disease | 0 |
| * Includes all positive HBsAg test results. | |
| Sexually Transmitted Disease | |
| Chlamydia Trachomatis | 32 |
| Gonorrhea | <5 |
| Syphilis | 0 |

IMMUNIZATIONS

| Children in Grades K-12 by Compliance Level | Number |
|---|--------|
| Compliant | 3,809 |
| Non-compliant | 5 |
| Percent Compliant | 99.9 |

Note: Death rates (except infant) are per 100,000 population. Infant, neonatal, and postneonatal death rates are per 1,000 live births. Perinatal and fetal death rates are per 1,000 live births plus fetal deaths. Rates are not calculated for fewer than 20 deaths.

| TOTAL DEATHS | | 223 |
|------------------|--------|--------|
| Crude Death Rate | | 885 |
| Age | Deaths | Rate |
| 1-4 | . | . |
| 5-14 | 2 | . |
| 15-19 | 2 | . |
| 20-34 | 2 | . |
| 35-54 | 29 | 407 |
| 55-64 | 28 | 700 |
| 65-74 | 45 | 1,659 |
| 75-84 | 59 | 3,829 |
| 85+ | 55 | 10,055 |

| Infant Mortality | | |
|------------------|--------|------|
| | Deaths | Rate |
| Total Infant | . | . |
| Neonatal | . | . |
| Postneonatal | . | . |

| Infant Mortality by Race of Mother | | |
|------------------------------------|--------|------|
| | Deaths | Rate |
| White | . | . |
| Black | . | . |
| Hispanic | . | . |
| Asian | . | . |
| Other/Unknown | . | . |

| Infant Mortality by Birthweight | | |
|---------------------------------|--------|------|
| | Deaths | Rate |
| <1,500 gm | . | . |
| 1,500-2,499 gm | . | . |
| 2,500+ gm | . | . |
| Unknown | . | . |

| Perinatal Mortality | | |
|---------------------|--------|------|
| | Deaths | Rate |
| Total Perinatal | . | . |
| Neonatal | . | . |
| Fetal | . | . |

| Selected Underlying Cause | Deaths | Rate |
|---------------------------|--------|------|
| Heart Disease (total) | 73 | 290 |
| Ischemic heart disease | 52 | 206 |
| Cancer (total) | 53 | 210 |
| Trachea/Bronchus/Lung | 15 | . |
| Colorectal | 5 | . |
| Female Breast | 2 | .* |
| Cerebrovascular Disease | 6 | . |
| Lower Resp. Disease | 9 | . |
| Pneumonia & Influenza | 2 | . |
| Accidents | 10 | . |
| Motor vehicle | 4 | . |
| Diabetes | 3 | . |
| Infect./Parasitic Dis. | 7 | . |
| Suicide | 7 | . |

* Based on female population.

ALCOHOL AND DRUG ABUSE AS UNDERLYING OR CONTRIBUTING CAUSE OF DEATH

| | Deaths | Rate |
|-------------|--------|------|
| Alcohol | . | . |
| Tobacco Use | 26 | 103 |
| Other Drugs | 2 | . |

MOTOR VEHICLE CRASHES

Note: These data are based on location of crash, not on residence.

| Type of Motor Vehicle Crash | Persons Injured | Persons Killed |
|-----------------------------|-----------------|----------------|
| Total Crashes | 129 | 6 |
| Alcohol-Related | 14 | 0 |
| With Citation: | | |
| For OWI | 17 | 0 |
| For Speeding | 24 | 0 |
| Motorcyclist | 12 | 0 |
| Bicyclist | 2 | 0 |
| Pedestrian | 0 | 0 |

2009 HOSPITALIZATIONS

| DISEASE / AGE GROUP | Number | Per 1,000 Pop | Average Stay (Days) | Average Charge | Charge Per Capita | DISEASE / AGE GROUP | Number | Per 1,000 Pop | Average Stay (Days) | Average Charge | Charge Per Capita |
|--|--------|---------------|---------------------|----------------|-------------------|---|--------|---------------|---------------------|----------------|-------------------|
| Injury-Related: All | | | | | | Alcohol-Related | | | | | |
| Total | 202 | 8.0 | 4.9 | \$26,434 | \$212 | Total | 29 | 1.2 | 3.8 | \$7,640 | \$9 |
| <18 | 11 | . | . | . | . | 18-44 | 14 | . | . | . | . |
| 18-44 | 41 | 5.3 | 4.0 | \$23,136 | \$122 | 45-64 | 14 | . | . | . | . |
| 45-64 | 56 | 6.9 | 4.9 | \$26,948 | \$185 | Pneumonia and Influenza | | | | | |
| 65+ | 94 | 19.6 | 5.4 | \$25,997 | \$509 | Total | 91 | 3.6 | 3.9 | \$12,000 | \$43 |
| Injury: Hip Fracture | | | | | | <18 | 6 | . | . | . | . |
| Total | 21 | 0.8 | 4.8 | \$26,420 | \$22 | 45-64 | 20 | 2.5 | 2.8 | \$10,215 | \$25 |
| 65+ | 20 | 4.2 | 4.8 | \$26,278 | \$109 | 65+ | 59 | 12.3 | 4.2 | \$11,847 | \$146 |
| Injury: Poisonings | | | | | | Cerebrovascular Disease | | | | | |
| Total | 17 | . | . | . | . | Total | 55 | 2.2 | 4.1 | \$27,793 | \$61 |
| 18-44 | 11 | . | . | . | . | 45-64 | 11 | . | . | . | . |
| Psychiatric | | | | | | 65+ | 42 | 8.7 | 3.4 | \$22,209 | \$194 |
| Total | 102 | 4.0 | 5.9 | \$8,109 | \$33 | Asthma | | | | | |
| <18 | 11 | . | . | . | . | Total | 10 | . | . | . | . |
| 18-44 | 55 | 7.1 | 5.3 | \$6,418 | \$45 | <18 | 0 | . | . | . | . |
| 45-64 | 28 | 3.4 | 5.7 | \$9,450 | \$32 | 18-44 | 3 | . | . | . | . |
| 65+ | 8 | . | . | . | . | 45-64 | 5 | . | . | . | . |
| Coronary Heart Disease | | | | | | 65+ | 2 | . | . | . | . |
| Total | 120 | 4.8 | 2.9 | \$37,764 | \$180 | Other Chronic Obstructive Pulmonary Disease | | | | | |
| 45-64 | 46 | 5.6 | 2.4 | \$33,698 | \$190 | Total | 53 | 2.1 | 4.8 | \$13,142 | \$28 |
| 65+ | 68 | 14.2 | 3.4 | \$41,095 | \$582 | 45-64 | 5 | . | . | . | . |
| Malignant Neoplasms (Cancers): All | | | | | | 65+ | 39 | 8.1 | 3.8 | \$10,178 | \$83 |
| Total | 100 | 4.0 | 5.4 | \$27,535 | \$109 | Drug-Related | | | | | |
| 18-44 | 2 | . | . | . | . | Total | 11 | . | . | . | . |
| 45-64 | 40 | 4.9 | 4.8 | \$23,015 | \$113 | 18-44 | 7 | . | . | . | . |
| 65+ | 57 | 11.9 | 5.8 | \$29,001 | \$344 | Total Hospitalizations | | | | | |
| Neoplasms: Female Breast (rates for female population) | | | | | | Total | 2,554 | 101.4 | 3.9 | \$19,120 | \$1,938 |
| Total | 4 | . | . | . | . | <18 | 303 | 67.7 | 3.7 | \$12,413 | \$841 |
| Neoplasms: Colo-rectal | | | | | | 18-44 | 523 | 67.4 | 3.5 | \$13,454 | \$906 |
| Total | 16 | . | . | . | . | 45-64 | 629 | 77.1 | 3.8 | \$22,533 | \$1,737 |
| 65+ | 10 | . | . | . | . | 65+ | 1,099 | 228.9 | 4.3 | \$21,713 | \$4,970 |
| Neoplasms: Lung | | | | | | ----- PREVENTABLE HOSPITALIZATIONS* ----- | | | | | |
| Total | 13 | . | . | . | . | Total | 361 | 14.3 | 3.8 | \$12,359 | \$177 |
| Diabetes | | | | | | <18 | 17 | 3.8 | 2.3 | \$7,545 | \$29 |
| Total | 24 | 1.0 | 3.0 | \$15,401 | \$15 | 18-44 | 37 | 4.8 | 5.5 | \$16,699 | \$80 |
| 65+ | 8 | . | . | . | . | 45-64 | 68 | 8.3 | 3.0 | \$10,143 | \$85 |
| | | | | | | 65+ | 239 | 49.8 | 3.9 | \$12,659 | \$630 |

* Hospitalizations for conditions where timely and effective ambulatory care can reduce the likelihood of hospitalization.