Wisconsin Family Health Survey

| Characteristics of Wisconsin's Household Population, 2016 | | | |
|---|---------|---------------|-----------|
| | Percent | 95% C.I. | Number |
| Total | 100.0% | | 5,622,000 |
| Age Groups | | | |
| 0 - 17 | 22 | (20.9 – 23.4) | 1,245,000 |
| 18 - 44 | 33 | (31.1 – 34.0) | 1,832,000 |
| 45 - 64 | 29 | (27.3 – 29.9) | 1,608,000 |
| 65 + | 17 | (15.7 – 17.7) | 937,000 |
| Sex and Age Groups | | | |
| Male | | | |
| 0 - 17 | 23 | (20.9 – 24.5) | 631,000 |
| 18 - 44 | 33 | (31.3 – 35.5) | 927,000 |
| 45 - 64 | 29 | (26.7 – 30.4) | 792,000 |
| 65 + | 15 | (14.0 – 16.7) | 426,000 |
| Female | | | |
| 0 - 17 | 22 | (19.8 – 23.3) | 614,000 |
| 18 - 44 | 32 | (29.7 – 33.8) | 905,000 |
| 45 - 64 | 29 | (26.9 – 30.5) | 817,000 |
| 65 + | 18 | (16.5 – 19.4) | 512,000 |
| Race/ethnicity | | | |
| White, non-Hispanic | 82 | (80.9 – 83.4) | 4,618,000 |
| African American, non-Hispanic | 6 | (5.0 – 6.5) | 325,000 |
| Hispanic/Latino | 6 | (5.4 – 7.2) | 357,000 |
| Asian, non-Hispanic | 1 | (1.1 – 1.7) | 78,000 |
| American Indian, non-Hispanic | 2 | (1.3 – 1.8) | 87,000 |
| Two or more races, non-Hispanic | 2 | (1.3 – 2.1) | 97,000 |
| Residence | | | |
| City of Milwaukee | 10 | (9.1 – 10.9) | 562,000 |
| Other Metropolitan (excluding | | | |
| city of Milwaukee) | 65 | (63.2 – 66.0) | 3,633,000 |
| Non-Metropolitan | 25 | (24.2 – 26.6) | 1,428,000 |
| Poverty Status | | | |
| Poor | 12 | (11.4 – 13.6) | 702,000 |
| Near-poor | 20 | (18.5 – 21.0) | 1,110,000 |
| Not poor | 67 | (65.4 – 68.3) | 3,757,000 |
| Educational Attainment | | | |
| Ages 18 and Older | | | |
| Less than high school diploma | 8 | (6.7 – 8.9) | 341,000 |
| High school graduate | 28 | (26.8 – 29.7) | 1,238,000 |
| Education beyond high school | 63 | (61.8 – 65.1) | 2,778,000 |

Wisconsin Family Health Survey

| Number 1,118,000 | | | |
|--|--|--|--|
| 1,118,000 | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| 127,000 | | | |
| | | | |
| 1,842,000 | | | |
| 170,00 | | | |
| | | | |
| 492,000 | | | |
| 929,000 | | | |
| Visconsin | | | |
| | | | |
| address-based design | | | |
| was used for sampling. However, comparisons with estimates from years of FHS data before 2012 | | | |
| should be made with caution because a landline telephone design was used for sampling prior to 2012. | | | |
| This table shows estimated percentages and numbers based on survey responses. C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical | | | |
| s). See Technical | | | |
| | | | |
| ent sample size to | | | |
| | | | |
| | | | |
| | | | |
| | | | |